


A11102 129258

NAT'L INST OF STANDARDS & TECH R.I.C.



A11102129258

/Scientific papers of the Bureau of Sten
QC1 .U572 V19:1923-24 C.1 NBS-PUB-C 1919







DEPARTMENT OF COMMERCE

SCIENTIFIC PAPERS

OF THE

BUREAU OF STANDARDS

GEORGE K. BURGESS, DIRECTOR

VOLUME 19

1923-24



WASHINGTON
GOVERNMENT PRINTING OFFICE
1925

30181

OCI

1312

7-

1810827
TO
GRADUATE
LIBRARY

CONTENTS OF VOLUME 19

	Page
469. DIRECTIVE RADIO TRANSMISSION ON A WAVE LENGTH OF 10 METERS. <i>Francis W. Dunmore and Francis H. Engel</i>	1
470. A METHOD FOR THE ACCURATE MEASUREMENT OF SHORT-TIME INTERVALS..... <i>Harvey L. Curtis and Robert C. Duncan</i>	17
471. METHODS OF MEASUREMENT OF PROPERTIES OF ELECTRICAL INSULATING MATERIALS..... <i>J. H. Dellinger and J. L. Preston</i>	39
472. ALTERNATING-CURRENT RESISTANCE AND INDUCTANCE OF SINGLE-LAYER COILS..... <i>C. N. Hickman</i>	73
473. A METHOD FOR THE MEASUREMENT OF SOUND INTENSITY... <i>J. C. Karcher</i>	105
474. SERIES IN THE ARC SPECTRUM OF MOLYBDENUM..... <i>C. C. Kiess</i>	113
475. VISIBILITY OF RADIANT ENERGY... <i>K. S. Gibson and E. P. T. Tyndall</i>	131
476. A STUDY OF RADIO SIGNAL FADING. <i>J. H. Dellinger, L. E. Whittemore, and S. Kruse</i>	193
477. SPECTRORADIOMETRIC ANALYSIS OF RADIO SIGNALS..... <i>Chester Snow</i>	231
478. REDETERMINATION OF SECONDARY STANDARDS OF WAVE LENGTH FROM THE NEW INTERNATIONAL IRON ARC. <i>W. F. Meggers, C. C. Kiess, and Keivin Burns</i>	263
479. INTERFEROMETER MEASUREMENTS OF THE LONGER WAVES IN THE IRON ARC SPECTRUM..... <i>W. F. Meggers and C. C. Kiess</i>	273
480. A DIRECTIVE TYPE OF RADIO BEACON AND ITS APPLICATION TO NAVIGATION..... <i>F. H. Engel and F. W. Dunmore</i>	281
481. MEASUREMENT OF LOW RESISTANCE BY MEANS OF THE WHEATSTONE BRIDGE..... <i>Frank Wenner and Alva Smith</i>	297
482. GRAVITATIONAL ANISOTROPY IN CRYSTALS..... <i>Paul R. Heyl</i>	307
483. INVESTIGATION OF THE PLATINUM METALS: IV. DETERMINATION OF IRIIDIUM IN PLATINUM ALLOYS BY THE METHOD OF FUSION WITH LEAD. <i>Raleigh Gilchrist</i>	325
484. PREPARATION AND PROPERTIES OF PURE IRON ALLOYS: IV. DETERMINATION OF THE CRITICAL RANGES OF PURE IRON-CARBON ALLOYS BY THE THERMOELECTRIC METHOD..... <i>J. F. T. Berliner</i>	347
485. APPLICATION OF THE INTERFEROMETER TO MEASUREMENTS OF THE THERMAL DILATATION OF CERAMIC MATERIALS..... <i>George E. Merritt</i>	357
486. SOME NEW THERMOELECTRICAL AND ACTINOELECTRICAL PROPERTIES OF MOLYBDENITE..... <i>W. W. Coblenz</i>	375
487. A QUANTITATIVE STUDY OF REGENERATION BY INDUCTIVE FEED BACK. <i>C. B. Jolliffe and Miss J. A. Rodman</i>	419
488. THERMAL EXPANSION OF MOLYBDENUM.... <i>Peter Hidnert and W. B. Gero</i>	429
489. PRIMARY RADIO-FREQUENCY STANDARDIZATION BY USE OF THE CATHODE-RAY OSCILLOGRAPH..... <i>Grace Hazen and Frieda Kenyon</i>	445
490. SPECTRA AND CRITICAL POTENTIALS OF FIFTH GROUP ELEMENTS. <i>Arthur E. Ruark, F. L. Mohler, Paul D. Foote, and R. L. Chenault</i>	463
491. THEORY OF DETERMINATION OF ULTRA-RADIO FREQUENCIES BY STANDARDING WAVES ON WIRES..... <i>August Hund</i>	487

	Page
492. FORMULAS, TABLES, AND CURVES FOR COMPUTING THE MUTUAL INDUCTANCE OF TWO COAXIAL CIRCLES. . . <i>Harvey L. Curtis and C. Matilda Sparks</i>	541
493. ULTRA-VIOLET REFLECTING POWER OF SOME METALS AND SULPHIDES. . . <i>W. W. Coblentz and C. W. Hughes</i>	577
494. ABERRATIONS OF LONG FOCUS ANASTIGMATIC PHOTOGRAPHIC OBJECTIVES . . . <i>A. H. Bennett</i>	587
495. A RADIOMETRIC INVESTIGATION OF THE GERMICIDAL ACTION OF ULTRA-VIOLET RADIATION. <i>W. W. Coblentz and H. R. Fulton</i>	641
496. EFFECT OF STRESS ON THE MAGNETIC PROPERTIES OF STEEL WIRE. <i>R. L. Sanford</i>	681
497. THERMAL EXPANSION OF ALUMINUM AND VARIOUS IMPORTANT ALUMINUM ALLOYS <i>Peter Hidnert</i>	697

DEPARTMENT OF COMMERCE
BUREAU OF STANDARDS
George K. Burgess, Director

SCIENTIFIC PAPERS OF THE BUREAU OF STANDARDS, No. 490
[Part of Vol. 19]

SPECTRA AND CRITICAL POTENTIALS OF
FIFTH GROUP ELEMENTS

BY

ARTHUR E. RUARK, Assistant Physicist
F. L. MOHLER, Physicist
PAUL D. FOOTE, Physicist
R. L. CHENAULT, Laboratory Assistant

Bureau of Standards

June 14, 1924



PRICE 10 CENTS

\$1.25 PER VOLUME ON SUBSCRIPTION

Sold only by the Superintendent of Documents, Government Printing Office
Washington, D. C.

WASHINGTON
GOVERNMENT PRINTING OFFICE
1924



SPECTRA AND CRITICAL POTENTIALS OF FIFTH GROUP ELEMENTS

By Arthur E. Ruark, F. L. Mohler, Paul D. Foote, and R. L. Chenault

ABSTRACT

Revised values of the critical potentials of As, Sb, and Bi are given. The resonance potentials of the neutral atoms correspond to the mean excitation voltages of groups of spectral lines. Photographs of the thermionic discharge in Bi were used to separate the excitation stages of the Bi atom; groups of spark lines appear at 14 ± 1 and 25 ± 5 volts. The Bi absorption spectrum was studied. Classifications of lines of N^+ , As, Sb, Bi, and Bi^+ were obtained. As, Sb, and Bi have doublet spectra. To describe phenomena connected with the existence of very narrow doublet levels in Al^+ , Bi, and other elements it is proposed to introduce a new quantum number called the fine quantum number.

CONTENTS

	Page
I. Introduction.....	463
II. Experimental work.....	465
1. Critical potentials.....	465
2. Absorption spectra.....	468
3. Spectroscopic study of stages in the thermionic discharge.....	469
III. Spectral classifications.....	472
1. Nitrogen.....	472
2. General characteristics of the spectra of As, Sb, and Bi.....	472
3. Impurity lines.....	474
4. Arsenic.....	475
5. Antimony.....	476
6. Bismuth.....	482
7. Higher stages in the bismuth spectra.....	485

I. INTRODUCTION

This paper discusses the arc spectra and critical potentials of arsenic, antimony, and bismuth, and includes some information on the spectrum of singly ionized nitrogen and the various spark spectra of bismuth. The data are voluminous, so we shall first outline the principal facts regarding these spectra. The elements nitrogen, phosphorus, arsenic, antimony, and bismuth lie on the right side of the fifth column of the periodic table. Their spectra are poor in lines compared with those of most elements on the right side of the periodic table. Stark and Hardtke¹ gave measures of a spectrum which they ascribe to the neutral

¹ Ann. der Physik., 56, p. 363; 1918.

atom of nitrogen, but the range of wave lengths is too narrow to justify an extended study of these lines. The great difficulty in obtaining the spectrum of neutral nitrogen is that the most common type of electron impact gives a molecule ion and an electron.² With low-current densities line spectra become visible only at about 70 volts.³ Probably a similar difficulty exists in the case of phosphorus. The line spectra of phosphorus have been studied by only a few observers, and the arc lines have not been well separated from the spark lines. Further observations will be required before this spectrum can be analyzed with economy of effort.

The conditions are quite different in the case of arsenic, antimony, and bismuth. As we pass to higher atomic weights in this column, dissociation can be obtained with weaker stimuli, so that the arc spectra of these three elements contain many reversed lines. Further, the lines due to the neutral atoms can be distinguished from the others with comparative ease. Thirty years ago Kayser and Runge⁴ discovered the most important constant differences in the arc spectra of these elements. Series of a conventional type were not found and no further classifications have been attempted since that time.

Our classification of the arc spectra of these elements shows that they possess doublet structure. The doublet spectra of the alkalis and of the aluminum group are known to be associated with the transitions of a single valence electron more loosely bound than any other electron in the atom. However, chemical evidence is not favorable to the assumption of a similar state of affairs in arsenic, antimony, and bismuth. Valences vary from 2 to 5, but are predominantly 3 and 5. It is the consensus of opinion that these three metals do not exhibit the valence one in any of their compounds. This strongly suggests that the arc spectrum arises from transitions of one electron of a group of three having azimuthal quantum numbers greater than unity.

We have studied the absorption spectra of arsenic, antimony, and bismuth, obtaining a line absorption spectrum only in the case of bismuth, and have also photographed the thermionic discharge in the latter element at potentials ranging between 4 and 60 volts. The spark lines of bismuth have been separated into at least three classes, the first of which appears at about 14 volts.

² Smyth, *Nature*, 111, p. 810; 1923.

⁴ *Ann. der Physik.*, 52, p. 93; 1894.

³ Duffendack, *Phys. Rev.*, 20, p. 665; 1922.

We have been able to classify nearly one-half the lines which can by any possibility belong to the neutral atoms of As, Sb, and Bi, and have found a fragmentary system of energy levels which accounts for some of the strongest lines in the 14-volt spectrum of Bi.

In As and Bi the critical potentials seem to be related to the atomic spectra in a simple manner, and have been used to obtain rough values for the spectroscopic terms. The question whether this procedure is justifiable in the case of Sb will be treated in the sequel.

We shall discuss, also, a multiplet found among the lines of N⁺.

II. EXPERIMENTAL WORK

1. CRITICAL POTENTIALS

Foote, Rognley, and Mohler determined the critical potentials of As, Sb, and Bi in 1918, but data were published only for As.⁵ The results for Sb and Bi were given in "The Origin of Spectra," without any experimental details and with cautions that they were not to be relied on. We have reexamined the curves for antimony and bismuth, and give slightly revised results for the whole family in Table 1. The values for antimony must still be considered as provisional.

TABLE 1.—Critical Potentials of the Nitrogen Group

Element	Inelastic collision potentials	Ionization
N.....*	8.18±0.1	16.9 ±0.5
P.....	5.80±0.1	13.3 ±0.5
As.....	4.69±0.1	11.54±0.5
Sb.....	{ 1.7 ±0.5 } 5.7 ±0.5	8.5 ±1.0
Bi ¹	1.9 ±0.1	8.0 ±0.5

¹ Inelastic collisions occur at 1.9, 3.9, and 5.7 volts. (See page 467.)

For these elements, the ionization potentials were observed by the method of total-current measurement and inelastic impacts were detected by the partial-current method in simple three-electrode tubes. In order to interpret the data for arsenic and antimony it is of importance to settle definitely the nature of the critical potentials for bismuth, these having been investigated in considerable detail. Temperatures between 700 and 800° C. were used for Bi. The measurements proved very difficult and

⁵ Phys. Rev., 13, p. 59; 1919.

gave rather discordant results. However, several of the better curves showed consistent values for the critical potentials which may be interpreted on the basis of the energy diagram (fig. 1).

Aside from the numerical agreements which are obtained, there is good evidence that these potentials are characteristic of the atom. The relative intensity of bands and lines in the absorption spectrum shows the presence of a large percentage of atoms at the temperatures of the low voltage arc experiments, as might be expected from the data on vapor pressure. The percentage would be increased by the dissociation due to the electron current.




FIG. 1.—Energy levels and prominent spectrum lines of arsenic, antimony, and bismuth

The extent of this increase is remarkable, for the low-voltage spectra are practically free of band structures. Those which do occur (in exposures taken at the beginning of the run) are not the bands appearing in the absorption spectrum, but are due to impurities.

Partial current measurements show prominently two drops in the current-voltage curves at an interval of 2.0 volts. Some curves show other drops, notably one at a potential 1.8 volts above the second resonance drop. From Figure 1 we see that atomic bismuth will have a large number of inelastic impact

potentials which will not be completely resolved by electrical methods. The three lowest, at 1.4, 1.9, and 2.6 volts, give rise only to weak spectral lines corresponding to zero change of the azimuthal quantum number, and are analogous to the metastable state potentials of the aluminum group. The mean value, 1.9 volts, corresponds well with the observed interval between the first two inelastic collisions. The raies ultimes 3067 and 4722 Å will be excited at 4.04 volts, so that the effect of this energy loss will be superposed on that of the second 1.9-volt loss. Thus we may use the first drop to determine the initial potential correction; it is found that the three impacts mentioned lie at 1.9, 3.9, and 5.7 volts. Possibly the drop at 5.7 is caused in part by combinations of the 1.9 and 4.0 volt losses, but it must be noted that many strong spectral lines, including the absorption lines 2276, 2230, 2228, 1959, and 1954 Å, will be excited at a mean potential of 5.8 volts. The ionization potential is 8.0 ± 0.5 volts, and this is in close agreement with the observed range of spectral terms. This measurement should be repeated, as the probable error is large, and the method used is probably not the best one for this element. We conclude that all these potentials are characteristic of the atom and that the lowest group of excitation potentials (mean value 1.9 volts) are the most prominent feature in the inelastic impact curves, although they give rise only to weak spectrum lines. In this respect the behavior of bismuth is quite similar to that of thallium.⁶

For arsenic and antimony also, there is little difficulty about obtaining a line spectrum free of bands, although the unexcited vapors at the temperatures used consist almost entirely of As_4 and Sb_4 . In arsenic at 400 to 500° C. it may be supposed that a considerable percentage of the atoms are in the states called $3D_1$, $3D_2$, and $3d_1$ (see fig. 1). The five lowest energy levels are so close together that the various inelastic collisions giving rise to transitions upward from these levels will not be separated in the electrical measurements. The observed potential is probably the superposed effect of the excitation voltages for a dozen or more spectrum lines between 3119 and 2349 Å. As a matter of fact, the mean voltage required to produce the seven raies ultimes (see Table 3) is 4.80, while the observed potential of inelastic impact is 4.7.

The experimental results for antimony are less reliable than those for bismuth and arsenic. No satisfactory correlation of

⁶ Mohler and Ruark, J. O. S. A. and R. S. I., 7, p. 819; 1923.

the critical potentials and the excitation stages of the arc spectrum has been attained, although the mean wave number of the raies ultimes agrees closely with the interval between the two observed potentials of inelastic collision. The energy diagram suggests the existence of other critical potentials much lower than this, in analogy to the 1.9-volt potential in Bi; however, it seems impossible to make a definite statement on the basis of existing data.

2. ABSORPTION SPECTRA

The absorption spectra of As, Sb, and Bi were photographed, utilizing the continuous background furnished by a spark under water, which passed between tungsten electrodes about 2.5 mm in diameter and 1 cm apart. A 1-kw. 20,000-volt radio transformer was connected to an oscillatory circuit containing a quenched spark gap, a glass plate condenser (immersed in oil) of about 0.005 microfarad, and the primary of a Tesla coil. This coil was made in accordance with a design described by Gibson and others;⁷ the secondary was connected direct to the terminals of the spark. A steady flow of distilled water was essential when exposures of more than a minute or two were required. If properly tuned this apparatus gives a background which does not contain any superposed emission or absorption lines. The limit of the spectrum is determined by the absorption bands of oxygen⁸ and by scattering due to the air and to the fused quartz plates in the path of the light. To record the lines 1954 and 1959 A of Bi an exposure of about 40 minutes, with the smallest size of Hilger quartz spectrograph, was required. For all other work a size E Hilger instrument was used, with exposures ranging between 2 and 40 minutes, depending on the spectral region to be examined. Satisfactory line absorption spectra of Bi were obtained at temperatures between 800 and 1,050° C. After this work was completed a paper by Grottrian⁹ appeared, dealing with the absorption spectra of several metals. He states that bismuth has an absorption line at 3067 Å and a group of bands in the region 2600 to 2900 Å, as well as a group in the visible region at higher temperatures. We have obtained absorption lines at 3067, 2276, 2230, 2228, 1959, and 1954 Å. 2276 Å is much stronger than 2230–2228 Å in absorption. Correspondingly it is usually observed to be weaker than 2230–2228 Å in emission, for it is reduced relatively more in intensity by the cooler vapors at the

⁷ B. S. Sci. Papers, 18, p. 128; 1922.

⁸ The first prominent oxygen band is at 1919.2 Å. Fainter ones occur at 1940 and 1957 Å. See L. and E. Bloch, C. R., 158, p. 1161; 1914.

⁹ Zeits. für Phys., 18, 169; 1923.

boundary of the luminous region. All the lines listed arise in transitions to the lowest level of the atom and are shown in the energy diagram of Bi (fig. 1). At $1,050^{\circ}$ C. there was not the slightest trace of spectrum lines arising on other low-lying orbits; indeed, the Boltzmann distribution law shows that they should not be expected.

At 800° C. scarcely a trace of bands could be observed, but at the higher temperatures and, accordingly, higher pressures, they were fully as prominent as the lines. A band group having about 17 heads lies between 2874 and 2672 Å. These bands are degraded toward the red. The frequency separation of corresponding points in neighboring bands is constant within the limits of our rough measurements, and amounts to about 175 cm^{-1} . Many of the bands have a narrow region of decreased absorption near their heads. A second group characterized by the same frequency separation begins at 2205 Å and extends toward shorter wave lengths. For this reason low temperatures (low pressures) had to be used in obtaining the lines 1959 and 1954 Å. Attempts were made to photograph line absorption spectra of arsenic and antimony. Both elements have a continuous absorption in the ultra-violet, which spreads toward longer wave lengths as the vapor pressure rises. The results of our experiments are in accord with the descriptions and photographs published by Dobbie and Fox.¹⁰

3. SPECTROSCOPIC STUDY OF STAGES IN THE THERMIONIC DISCHARGE

Bismuth was boiled in a silica tube sealed at the bottom and closed at the top with a water-cooled metal plate. A tube carrying the electrodes and the observing tube passed through holes in the metal top.

The tungsten cathode was surrounded by a cylindrical sheet steel anode about 1 cm in diameter and the inside of the quartz tube was lined with sheet steel which was kept at the same potential as the anode. The lower end of the tube was heated by an electric furnace to temperatures between 800 and 900° C. Outside the furnace and tube was mounted a coil of copper wire. A large current in this coil maintained a magnetic field, which concentrated the discharge in a beam, nearly parallel to the line of sight.

Thermionic currents up to 1 ampere could be maintained above the ionization potential, but at lower voltages 10 milliamperes was about the maximum current. Light reflected diffusely from the incandescent cathode was troublesome and set a limit (about two

¹⁰ Proc. Roy. Soc. A., 98, p. 147; 1920.

hours) to the length of exposure that was useful in photographing the low-voltage stages. For this reason the work with pure bismuth was supplemented with a few photographs of discharges in a mixture of sodium and bismuth. Large currents could then be maintained at potentials slightly above the ionization potential of sodium, 5.1 volts. This made it possible to determine excitation stages for faint lines and lines in the far ultra-violet.

In estimating the true potential in pure bismuth it was assumed that the potential at the arc point was 8 volts. This gave a correction to the applied potential ranging between +0.5 and +0.8, but the values obtained by this method are subject to errors of possibly 1 volt.

In discharges above the arcing potential mean-current densities of the order of 0.1 to 1 amperes per cm^2 were used.

The spectral region studied extends from 4722 to 2157 Å. Exposures were obtained from 4 to 60 volts; those taken below 8 volts confirm the classification of the lines of neutral Bi. As nearly as could be judged, the more prominent lines appeared in every case at voltages agreeing with those predicted from the classification. A 5-volt exposure showing only the resonance lines 3067 and 4722 Å is reproduced in Figure 2. The 8-volt exposure is characteristic of the arc spectrum, while the 16-volt photograph shows the first set of lines to appear after the arc spectrum. At 60 volts nearly all the spark lines recorded by more than one observer are present.

The photographs taken above the ionization potential include the following voltages: 10, 14, 15, 16, 20, 25, 30, 35, 40, 45, 50, and 60. Table 2 gives a list of the spark lines observed. Opposite each line is tabulated the voltage at which it first appears upon our plates, with remarks in some cases as to the manner of enhancement with increasing voltage.

Schuster and Hemsalech¹¹ have studied the velocities with which the luminous vapors in the bismuth spark pass outward from the poles, while Hemsalech¹² has described the effects of self-induction on prominent Bi lines. Their results afford support to the assignment of the excitation stages for the more prominent lines in Table 2, and enable us to state, in addition, that 5209 and 5271 Å belong to a higher stage than the 14-volt lines. In attempting to obtain from Table 2 the approximate critical potentials of the spark spectra, the question arises whether excitation by electron collisions with ions was probable. High-current

¹¹ Phil. Trans. A., 193, p. 189; 1900.

¹² Jour. de Physique (3), 8, p. 652; 1899.




FIG. 2.—Absorption spectrum and stages in the excitation of bismuth

densities favor this type of excitation, and by studying the discharge with different current densities at each potential used, it should be possible to determine the origins of many spectral lines and of the critical potentials encountered. Such a study was not made for Bi, but comparison with results obtained with other elements indicates that collisions of electrons with ions will be an appreciable factor at the current densities used.

The first stage in the spark spectrum appears at 14 ± 1 volts. Other stages occur at 25 ± 5 and probably at 45 ± 5 volts. Intermediate stages may occur, as the excitation potentials of the many fainter lines are very uncertain.

Spectroscopic stages in arsenic and antimony have not been studied in detail, although photographs of the antimony spectrum have been obtained at several potentials up to 30 volts.

TABLE 2.—Appearance Voltages of Bi Lines

λ I. A.	Intensity	Voltage of appearance	Remarks and classification
2,297.5	60	
2,345.9	2U	50	
2,368.40	2	16	
2,414.80	20	20	On one plate only. May be Hg.
2,499.51	1	25	
2,500.5	35	
2,544.4	1u	14	Much enhanced at higher voltages.
2,594.05	1	8.5	
2,679.9	30	
2,692.9	15	
2,713.4	2u	16	Do.
2,746.35	1	25	
2,803.48	4	14	} Not separated. Enhanced at higher voltages.
2,803.69	5	14	
2,805.2	14	
2,847.6	1u	50	Enhanced at higher voltages.
2,855.68	30	30	
3,111.4	1u	50	
3,115.4	2u	14	
3,187.6	15	Visible at 15, but strongly enhanced at higher voltages.
3,296.0	14	
3,299.7	1u	14	G—N.
3,315.2	14	
3,393.1	1u	14	} Not separated on our plates.
3,394.1	1u	14	
3,431.01	20	14	G—L.
3,451.05	20	45	
3,455.2	1u	14	G—K. Wave length adjusted.
3,473.9	5u	40	
3,485.6	5u	14	
3,541.4	2u	35	Weak at 35 volts.
3,613.8	5r	25	
3,654.4	1u	14	H—N.
3,695.55	50	25	Weak at 25 volts.
3,756.4	2u	14	
3,792.8	50u	14	
3,811.2	2u	14	
3,816.1	4u	14	H—L.
3,846.0	2u	25	H—K. Possibly present at 20 volts.
3,864.2	10u	14	
4,079.25	30	14	
4,259.70	100	14	
4,302.10	50	14	
4,340.6	2u	14	Probably due to lead.
4,386.8	2u	8.5	Not recorded in the arc in air.
4,391.4	1u	16	
4,477.1	1u	25	Faint at all voltages.
4,561.15	20R	45	

III. SPECTRAL CLASSIFICATIONS¹³

1. NITROGEN

Professor Fowler¹⁴ has pointed out seven lines almost certainly due to doubly ionized nitrogen, which exhibit the doublet regularities characteristic of the neutral atoms of third group elements. We have found a six-line multiplet of the type $mp-np$, which obeys the intensity rules fairly well. These lines are probably due to singly-ionized nitrogen, for they appear at a lower excitation than Fowler's doublets. This striking group is shown in Plate XIII of Fowler's article. The data are as follows:

λ I. A.	Intensity	ν	Classification
4,643.10.....	7	21,531.29	mp_1-np_2
4,630.55.....	10	21,589.67	mp_1-np_1
4,621.40.....	7	21,632.40	mp_2-np_3
4,613.88.....	6	21,667.65	mp_2-np_2
4,607.16.....	7	21,699.24	mp_3-np_2
4,601.48.....	8	21,726.02	mp_2-np_1

Attention may also be drawn to the frequency difference, 51.69, occurring among the lines of singly-ionized nitrogen:

4,236.99	4 h	23,595.01
4,227.74	2 h	23,646.62
4,043.44	2 h	24,723.84
4,035.09	4 h	24,775.59

2. GENERAL CHARACTERISTICS OF THE SPECTRA OF As, Sb, AND Bi

The arc spectrum of arsenic is very poor in lines. There are no lines in the visible region, for two reasons: (1) There is a gap of 32,000 cm^{-1} in the energy diagram, lying just above the five lowest orbits, and (2) the terms lying above the gap do not combine in such a way as to yield lines in this region, although many such combinations are numerically possible. A similar phenome-

¹³ The following abbreviations are used to indicate the character of spectrum lines:

R=reversed.
u or h=hazy, diffuse.
U or H=very diffuse.
b=broad.
B=very broad.
r=diffuse toward red.
v=diffuse toward violet.

In this paper all wave lengths recorded originally on the Rowland scale have been reduced to I. A., with the exception of the infra-red measures of Randall. For these the errors estimated by Randall are larger than the corrections, and the latter are not known with any certainty beyond 8800 Å. The wave numbers given are in every case the reciprocals of the wave lengths in vacuo obtained from the wave lengths listed in the tables. The corrections to vacuum were made in accordance with the data of Meggers and Peters, B. S. Bull. 14, p. 697; 1918. The series notation is that of Paschen.

¹⁴ Monthly Notices, R. A. S., 80, 692; 1920.

non is found in Sb and Bi. The gap in the spectrum becomes narrower and shifts toward longer wave lengths as we pass to higher atomic numbers. In each case the vacant region in the energy diagram is just above a comparatively compact group of low-lying orbits (see fig. 1). In As and Sb this group consists of five orbits. From the nature of the combinations we have designated these orbits in arsenic as $3D_1$, $3D_2$, $1s$, $3d_1$, and $3d_2$. The capital letters are used only to avoid some distinguishing mark, such as a prime, because it seems reasonable to assign the same term number to both $3D$ and $3d$ orbits. In all probability these pairs of orbits are characterized by the same total, azimuthal and inner quantum numbers, the difference of energy being attributable to screening. It will be understood that the orbits $4d_{1,2}$ might just as well be designated as $4D_{1,2}$.

In Bi the separations of the levels corresponding to the $3d$ and $3D$ doublets in As are, of course, very large, and a new feature appears. In place of the energy level $3D_1$ of As we have a pair of energy levels separated by 1.85 cm^{-1} , and in place of the energy level $3d_1$ we have a pair separated by 14.05 cm^{-1} . (Compare, however, the alternative classifications mentioned on page 483.)

By analogy with arsenic, the first pair is designated as $3D_1^A$ and $3D_1^B$, the second, as $3d_1^A$ and $3d_1^B$, while the remaining levels of the group of seven are assigned the same names as the arsenic levels which occupy corresponding positions. The group of five in Sb is given the same notation as in arsenic, but these assignments may have no physical significance. The combinations in the Sb and Bi spectra are such that we are not justified in specifying the character of their energy levels at the present stage of our knowledge.

The 1.85 cm^{-1} separation in Bi gives rise to extremely narrow doublets in the visible and ultra-violet regions of the spectra, and it is not to be expected that the corresponding doublets in As and Sb could be resolved except by a very laborious and elaborate study. However, the analogues of the 14 cm^{-1} separation should be observed with reasonable ease, and it would be of considerable interest to determine whether they actually exist. A somewhat similar term multiplicity was observed by Paschen¹⁵ in the spectrum of Al^+ , which consists of singlet and triplet series. Paschen found that the terms $4f_1$, $4f_2$, $4f_3$, and $4F$ are each very narrow doublets. From a study of fine structures observed in the spectra of Cu, Hg, Al^+ , Tl, and Bi, we conclude that such mul-

¹⁵ Ann. d. Physik., **71**, p. 537; 1923.

tiplicities require the introduction of a new quantum number, for which we propose the name "fine quantum number" and the designation f . It is reasonable to suppose that small quantized differences in the energy of the interior electron shells (or perhaps different orientations of an asymmetric nucleus) correspond to the various values of the fine quantum number. While there may be fine structures in the line spectra of isotopic atoms which are brought about by the variations in the mass or form of the nucleus, we have good evidence, based on the combinations observed in the elements mentioned above, that there are other types of fine structure which arise entirely from variations in the energy levels of a single atom. The facts may be illustrated by a simple case. If we observe the spectrum of an element of given atomic number, which is composed of two isotopes, we should expect that each energy level would appear to be split into two levels, which we may designate as "low" and "high." Combinations between "low" levels and "high" levels could not occur, while cases are known in which three or four fine structure components arise from two energy levels, each consisting of a narrow doublet. It was through the study of the fine structures observed in Bi that we were led to classify these spectra.

The magnitudes of the highest terms in these spectra were estimated from the resonance and ionization potentials. It is believed the assignments in As and Bi are not in error by more than $4,000\text{ cm}^{-1}$. The critical potentials of Sb have not been correlated with the spectrum in a satisfactory manner, and the uncertainty of the term magnitude is still very great.

The constant differences discovered by Kayser and Runge were a great aid in finding further regularities. They pointed out the relationships of 20 lines in As, 34 in Sb, and 25 in Bi. All of the differences noted were genuine with the exception of two in Sb. This changes the classification of 2395, 2212, 2352, and 2175 Å.

3. IMPURITY LINES

No extensive attempt was made to locate impurity lines in the lists of spark wave lengths in Kayser's Handbuch, but the following were noted:

In the Arsenic Lists.

Bi: 4722.62, 4259.61, 3067.82.

Pb: 4057.94, 3740.12, 3683.56, 3671.90, 3639.74, 3572.94, 2833.2, 2802.1, 2663.22, 2614.09.

Sb: 4352.28?, 3267.60, 3232.67, 2598.10, 2528.49, 2445.61, 2306.7?.

In the Antimony Lists

Al: 3961.63, 3944.16.
As: 5651.39, 5558.06, 5497.73, 4552.29, 4543.64, 4539.84, 4494.48, 4474.44, 4466.48, 4458.61, 4368.90, 4336.77, 4249.61?, 4082.60, 4006.21, 3948.69, 2349.84, 2288.21.
Bi: 3024.85?.
Cu: 5782.15, 5646.3, 5599.76, 5463.7, 5218.14, 3273.97, 3247.55.
Pb: 5609.2, 5372.9, 4387.01, 4245.34, 4058.00, 3739.95, 3683.47, 3639.57, 2663.31, 2614.33, 2246.99, 2203.65, 2170.23.
Si: 2528.58, 2519.31, 2218.15, 2216.52?.
Sn: 2706.73, 2571.64, 2429.55.

Walters has pointed out that several of the lines given by Kretzer between 7145 and 7480 Å are second order lines due to Pb and Sb. Two lines were outstanding with wave lengths 7343.41 and 7181.09 Å. The first is Pb 3671 Å, and there can be little doubt that the sole survivor is also spurious. Kretzer's line at 6495.17 may be the second order of Cu 3247.66 Å.

In the Bismuth Lists

Ag: 3381.9, 3279.9, 2437.2.
Cd: 2265.11, 2144.58?.
Hg: 2847.7?, 2224.73?.
Pb: 4387.0?, 3740.10, 3683.64, 3639.75, 3572.92, 2613.74, 2446.3, 2402.08, 2394.0.

The following lines listed by Kayser and Runge as occurring in the arc did not appear on our plates: 2944.38, 2892.98, 2582.17, 2532.65, 2489.5, 2360.0, 2309.4, 2281.39, 2224.27, and 2203.2 Å.

4. ARSENIC

The Zeeman effect of arsenic has not been studied, and no measures have been made from 8823 to 6404 Å. Randall¹⁶ gives wave lengths from 10614 to 8823 Å. We do not give the list, since no classifications of any certainty were obtained. The characteristic arc spectrum extends from 3119 Å toward shorter wave lengths. Kayser's and Runge's measurements were used to 2144 Å, inclusive, and those of L. and E. Bloch¹⁷ beyond that point. The wave lengths are recorded only to 1831 Å, since we could not classify any shorter radiations. Tables 3 and 4 give the classifications and term magnitudes, respectively, while Table 5 shows the nature of the combinations. The assignment of inner quantum numbers j depends on the fact that the lines $3d_1-2p_2$ and $3D_1-2p_2$ are absent. If we admit that this is an instance of the selection principle for inner numbers, the values adopted are necessary (of course, they can all be changed

¹⁶ *Astrophys. J.*, **34**, p. 1; 1911.

¹⁷ *Jour. de Physique*, **4**, p. 622, 1914; and *C. R.*, **171**, p. 709; 1920.

by a constant), and the intensity rules are then found to be fairly well obeyed.

The following pairs are of doubtful significance:

λ	ν	
8,823.3	11,330.6	} $\Delta\nu = 916.0$; $\Delta_3 D = 916.23$
9,599.3	10,414.6	
1,847.7	54,121	} $\Delta\nu = 830$; $\epsilon - \zeta = 829.68$
1,872.28	53,392	

A convenient point of attack for the discovery of further regularities may be afforded by the narrow pairs at wave lengths 1741.6, 1739.4, and 1616.7, 1614.8 Å, which exhibit the difference 72.6 cm^{-1} . The pairs 1634.0, 1630.9 and 1574.7, 1571.7 Å are also worthy of attention.

5. ANTIMONY

The wave lengths of Randall (*loc. cit.*) extend from 12118 to 9519 Å. In the region 9132 to 5531 Å we have used Walters'¹⁸ data, supplemented where necessary by those of Schippers and of Kretzer, as listed in Kayser's Handbuch. Walters found that some lines were very much stronger near the positive electrode than in any other part of the arc. These lines are indicated by the sign + in the intensity column of Table 7; together with a few other lines they have been designated as spark lines in the column of remarks. From 5490 to 2068 Å, the wave lengths are those of Schippers, while the data of L. and E. Bloch¹⁹ were used from 2063 to 1949 Å. Beyond that point we have adjusted the wave lengths of Takamine and Nitta,²⁰ McLennan, Young, and Ireton,²¹ and L. and E. Bloch.²² Table 6 gives the terms of the antimony spectrum, while Table 7 contains wave lengths and classifications. The Zeeman effect in antimony and bismuth was studied by Purvis²³ and by Van der Harst,²⁴ but the results have not been of any aid in classification.

It may be noted that the assumption of an energy level E at 25669.41 cm^{-1} gives the following lines:

$$3d_2 - E = 45,184,$$

$$E - \kappa = 10,047.64 \text{ while } 10,047.02 \text{ is observed.}$$

$$E - \nu_2 = 11,513.31 \text{ while } 11,513.76 \text{ is observed.}$$

Since these classifications are uncertain they are not included in Table 7.

¹⁸ B. S. Sci. Papers, 17, p. 161; 1921.

¹⁹ Jour. de Physique, 4, p. 622; 1914.

²⁰ Kyoto Coll. of Sci. Memoirs, 2, p. 117; 1917.

²¹ Proc. Roy. Soc. 98, p. 95; 1920.

²² C. R., 171, p. 709; 1920.

²³ Proc. Cambridge Phil. Soc., 14, p. 216; 1907.

²⁴ Amsterdam Proceedings, 22, p. 300; 1920.

TABLE 3.—Term Magnitude in As

Name of term	j	Term magnitude	Difference
θ	10,383.86	1,505.45
η	11,889.31	689.10
ζ	12,578.41	829.68
ϵ	13,408.09	477.62
δ	13,885.71
γ	40,550.96
$2p_1$	2	51,147.28
$2p_2$	1	51,469.61	322.33
$4d_1$	3	58,741.04
$4d_2$	2	59,202.18	461.14
$3D_1$	3	91,248.90
$3D_2$	2	92,165.13	916.23
$1s$	93,452.43
$3d_1$	3	93,690.27
$3d_2$	2	95,160.00	1,469.73

TABLE 4.—Arc Spectrum of Arsenic

λ	Intensity	ν	Classification	Remarks
λ vac				
1,831.1.....	1	54,612	$3d_2-\gamma$	
1,844.3.....	3	54,221	
1,847.7.....	2	54,121	
1,850.6.....	2	54,037	
1,854.4.....	2	53,926	
λ I. A.				
1,859.90.....	1	53,748	Extrapolated.
1,871.07.....	1	53,427.4	
1,872.33.....	1	53,391.4	
1,881.08.....	1	53,143.2	$3d_1-\gamma$	
1,889.91.....	4R	52,894.9	$1s-\gamma$	
1,912.27.....	1	52,275.4	
1,916.48.....	1	52,161.6	
1,936.94.....	5R	51,610.8	$3D_2-\gamma$	
1,958.31.....	2	51,021.7	
1,971.97.....	4R	50,694.0	$3D_1-\gamma$	
1,990.21.....	2	50,229.6	
1,994.74.....	2	50,115.5	Diffuse.
2,003.35.....	2	49,900.2	Do.
2,009.18.....	2	49,755.4	Do.
2,010.10.....	49,732.7	Kayser and Runge's λ , adjusted
2,013.11.....	2	49,658.4	Diffuse.
2,047.75.....	2	48,818.4	$4d_2-\theta$	
2,065.43.....	2	48,400.6	
2,067.29.....	1	48,357.1	$4d_1-\theta$	
2,068.43.....	1	48,330.4	
2,069.87.....	2	48,296.9	
2,079.41.....	1	48,075.1	Coincident with a line of Cu.
2,085.27.....	2	47,940.0	Kayser and Runge's λ , adjusted.
2,089.04.....	47,853.5	
2,089.73.....	1	47,837.7	Diffuse.
2,095.10.....	2	47,715.2	
2,109.80.....	1	47,382.8	
2,112.93.....	3	47,312.6	$4d_2-\eta$	
2,133.70.....	2	46,852.0	$4d_1-\eta$	
2,144.15.....	4	46,623.8	$4d_2-\zeta$	
2,165.57.....	4	46,162.6	$4d_1-\zeta$	
2,176.30.....	1	45,935.1	
2,183.00.....	1	45,794.0	$4d_2-\epsilon$	
2,205.21.....	2	45,333.0	$4d_1-\epsilon$	
2,206.01.....	2	45,316.5	$4d_2-\delta$	
2,228.70.....	2	44,855.3	$4d_1-\delta$	
2,266.72.....	4	44,103.0	
2,271.39.....	4	44,012.4	$3d_2-2p_1$	
2,288.12.....	10R	43,690.5	$3d_2-2p_2$	Raie ultime.
2,349.84.....	10R	42,543.0	$3d_1-2p_1$	Do.

TABLE 4.—Are Spectrum of Arsenic—Continued

λ	Intensity	ν	Classification	Remarks
λ I. A.				
2,363.04.....	2	42,305.4	$1s-2p_1$	
2,369.67.....	4R	42,187.1	
2,370.77.....	4R	42,167.4	
2,381.20.....	4R	41,982.7	$1s-2p_2$	
2,437.22.....	1	41,018.0	$3D_2-2p_1$..	
2,456.53.....	4R	40,695.6	$3D_2-2p_2$..	
2,492.89.....	1	40,102.0	$3D_1-2p_1$..	
2,744.99.....	6R	36,419.2	$3d_2-4d_1$..	Raie ultime.
2,780.20.....	8R	35,958.0	$3d_2-4d_2$..	Do.
2,860.43.....	6R	34,949.5	$3d_1-4d_1$..	Do.
2,898.72.....	4R	34,487.9	$3d_1-4d_2$..	Do.
2,990.99.....	2	33,424.0	$3D_2-4d_1$..	
3,032.84.....	4	32,962.8	$3D_2-4d_2$..	
3,075.32.....	2	32,507.6	$3D_1-4d_1$..	
3,119.57.....	4	32,046.5	$3D_1-4d_2$..	

TABLE 5.—Combinations Occurring in the As Spectrum

[A cross indicates that the line is present. Rales ultimes are distinguished by stars]

	$4d_2$	$4d_1$	$2p_2$	$2p_1$	γ	δ	ϵ	ζ	η	θ
$3d_2$	×	×	×	×	×					
$3d_1$	×	×	×	×	×					
$1s$			×	×	×					
$3D_2$	×	×	×	×	×					
$3D_1$	×	×		×	×					
$4d_2$						×	×	×	×	×
$4d_1$						×	×	×	×	×

TABLE 6.—Term Magnitudes in Antimony

Name of term	Term magnitude	Differences
ω	5,881.56	
ψ	9,047.48	
χ	9,468.92	
φ	9,854.16	
ν	10,274.31	
τ	10,450.3	
σ	11,991.63	
ρ	12,019.05	
π	12,108.18	
ϕ_1	12,201.47	
ϕ_2	12,265.02	63.55
ξ	13,257.70	
ν_1	14,121.61	
ν_2	14,156.10	34.49
μ	14,702.58	
λ	15,125.74	
κ	15,621.77	
ϵ_1	15,719.97	
ϵ_2	15,734.58	14.61
θ	17,326.72	
η	17,411.54	
ζ	19,770.86	
ϵ	21,463.37	
δ	22,521.60	
γ	23,863.41	
β	24,909.00	
α	27,604.98	
$3D_1$	52,390.28	
$3D_2$	54,459.05	2,068.77
$1s$	61,000.00	
$3d_1$	62,341.61	
$3d_2$	70,853.47	8,511.86

TABLE 7.—Arc Spectrum of Antimony

λ	Intensity	ν	Classification	Remarks
λ Vac.				
1,437.0.....	7	69,589	
1,438.7.....	4	69,507	
1,493.3.....	4	66,966	
1,495.6.....	4	66,863	
1,499.25....	7	66,700	
1,504.85....	6	66,552	
1,506.4.....	66,383	
1,514.1.....	7	66,046	
1,533.5.....	4	65,210	
1,535.8.....	2	65,113	
1,540.7.....	2	64,906	
1,554.65....	2	64,323	
1,566.2.....	63,849	
1,574.9.....	2	63,496	
1,585.3.....	63,079	
1,600.8.....	62,469	
1,607.6.....	4	62,204	
1,613.9.....	1	61,962	
1,631.5.....	2	61,293	
1,635.0.....	61,162	
1,640.8.....	3	60,946	
1,667.2.....	59,981	
1,674.2.....	59,730	
1,678.1.....	59,591	$3d_2-\nu$	
1,699.0.....	4	58,858	$3d_2-\rho$, or $3d_2-\sigma$	
1,702.3.....	4	58,744	$3d_2-\pi$	
1,712.0.....	58,411	
1,717.1.....	58,238	
1,725.3.....	57,961	
1,730.7.....	57,780	
1,735.9.....	5	57,607	$3d_2-\xi$	
1,762.45....	10	56,739	$3d_2-\nu_1$, or $3d_2-\nu_2$	
1,780.55....	4	56,162	$3d_2-\mu$	
1,788.05....	2	55,927	
1,799.95....	5	55,557	
1,810.15....	5	55,244	$3d_2-\kappa$	
1,814.25....	6	55,119	$\{55,133=3d_2-\iota_1$ $55,118=3d_2-\iota_2$ or $1s-\omega$
1,817.6.....	55,018	
1,821.3.....	54,906	
1,829.4.....	5	54,663	
1,832.8.....	54,561	
1,839.1.....	54,374	
1,867.8.....	8	53,539	$3d_2-\theta$	
1,870.75....	10	53,454.5	$3d_2-\eta$	
1,877.91....	53,250.7	
1,882.45....	4	53,122.3	
1,891.05....	3	52,908.6	
1,899.45....	3	52,646.8	
1,922.68....	52,010.7	
1,926.61....	5	51,904.6	
1,930.83....	5	51,791.2	
λ I. A.				
1,949.58....	5	51,276.3	
1,977.58....	3	50,550.2	$3d_1-\tau$	
1,985.30....	3	50,352.2	$3d_1-\sigma$	
2,004.77....	3	49,864.9	
2,014.69....	2	49,619.4	
2,023.86....	4	49,394.7	$3d_2-\epsilon$	
2,029.27....	4	49,262.8	
2,036.51....	2	49,087.7	$3d_1-\xi$	
2,039.60....	5	49,013.4	$1s-\sigma$	
2,044.56....	4	48,894.5	$1s-\pi$	
2,046.48....	3	48,848.6	
2,049.46....	7	48,777.6	Diffuse.
2,054.04....	6	48,668.9	Probably a spark line.
2,061.92....	3	48,483.0	
2,063.44....	5	48,447.3	Diffuse.

TABLE 7.—Arc Spectrum of Antimony—Continued

λ	Intensity	ν	Classification	Remarks
λ I. A.				
2,068.38....	2R	48,331.6	$3d_2-\delta$	
2,079.63....	1	48,070.0	
2,098.47....	2	47,638.6	$3d_1-\mu$	
2,117.28....	2	47,215.5	$3d_1-\lambda$	
2,127.46....	3	46,989.6	$3d_2-\gamma$	
2,137.11....	3	46,777.3	
2,139.76....	3R	46,719.4	$3d_1-\kappa$	
2,141.76....	3	46,675.7	
2,145.04....	3	46,604.4	$3d_1-\epsilon_2$	
2,158.97....	4	46,303.8	
2,159.26....	3	46,297.6	$1s-\mu$	Kayser and Runge.
2,175.88....	5R	45,944.0	$3d_2-\beta$	
2,179.26....	4R	45,872.6	$1s-\lambda$	
2,201.39....	3R	45,411.6	$3D_1-\psi$	
2,203.03....	2	45,377.8	$1s-\kappa$	
2,207.79....	2	45,280.0	$1s-\epsilon_1$	Do.
2,208.52....	3R	45,265.0	$1s-\epsilon_2$	
2,212.48....	1	45,184.1	
2,220.81....	2	45,014.6	$3d_1-\theta$	
2,222.05....	2	44,989.5	$3D_2-\chi$	
2,225.00....	2	44,929.9	$3d_1-\eta$	
2,262.54....	3	44,184.4	$3D_2-\nu$	
2,289.01....	3	43,673.5	$1s-\theta$	
2,293.47....	4	43,588.6	$1s-\eta$	
2,306.48....	4R	43,342.8	$3D_1-\psi$	
2,311.50....	6R	43,248.7	$3d_2-\alpha$	Raie ultime
2,329.09....	2	42,922.0	$3D_1-\chi$	
2,352.20....	2	42,500.4	
2,360.50....	3	42,351.0	$3D_2-\pi$	
2,373.65....	4R	42,116.3	$3D_1-\nu$	
2,383.63....	4R	41,940.0	$3D_1-\tau$	
2,395.22....	2	41,737.1	
2,422.15....	3	41,273.0	
2,426.36....	4R	41,201.4	$3D_2-\xi$	
2,445.53....	4R	40,878.6	$3d_1-\epsilon$	
2,474.58....	4	40,398.6	$3D_1-\sigma$	
2,478.34....	2	40,337.4	$3D_2-\nu_1$	
2,480.45....	3	40,303.1	$3D_2-\nu_2$	
2,481.75....	3	40,282.0	$3D_1-\pi$	
2,510.54....	3	39,820.0	$3d_1-\delta$	
2,514.56....	2	39,756.4	$3D_2-\mu$	Do.
2,528.54....	6R	39,536.6	$1s-\epsilon$	
2,554.65....	3	39,132.5	$3D_1-\xi$	
2,574.09....	4	38,837.1	$3D_2-\kappa$	
2,598.08....	6R	38,478.4	$\left\{ \begin{array}{l} 3d_1-\gamma \\ \text{or } 1s-\delta \end{array} \right.$	$\left\{ \begin{array}{l} \text{Do.} \\ 3d_1-\gamma=38,478.20, \\ 1s-\delta=38,478.40. \end{array} \right.$
2,612.32....	4R	38,268.7	$3D_1-\nu_1$	
2,614.69....	2	38,234.0	$3D_1-\nu_2$	
2,652.61....	4R	37,687.5	$3D_1-\mu$	
2,670.66....	5R	37,432.8	$3d_1-\beta$	
2,682.76....	4R	37,264.9	$3D_1-\lambda$	
2,692.27....	3R	37,132.3	$3D_2-\theta$	
2,718.91....	4R	36,768.6	$3D_1-\kappa$	
2,727.22....	3	36,656.5	$3D_1-\epsilon_2$	
2,769.94....	5R	36,091.2	$1s-\beta$	
2,851.12....	2	35,063.6	$3D_1-\theta$	
2,877.92....	5R	34,737.2	$3d_1-\alpha$	Raie ultime.
3,029.82....	4R	32,995.7	$3D_2-\epsilon$	
3,232.54....	4R	30,926.5	$3D_1-\epsilon$	Do.
3,267.50....	4R	30,595.6	$3D_2-\gamma$	
3,383.15....	3	29,549.8	$3D_2-\beta$	
3,504.45....	2	28,527.0	$3D_1-\gamma$	
3,637.83....	3	27,481.1	$3D_1-\beta$	
3,722.79....	3	26,854.0	$3D_2-\alpha$	
4,033.56....	3	24,785.0	$3D_1-\alpha$	
5,490.33....	1	18,208.80	

TABLE 7.—Arc Spectrum of Antimony—Continued

λ	Intensity	ν	Classification	Remarks
λ I. A.				
5,531.89.....	1, h	18,072.01	-----	Spark line.
5,556.21.....	3	17,992.89	-----	
5,568.09.....	3	-----	-----	
5,599.93.....	3	17,852.42	-----	
5,602.30.....	3	17,844.87	-----	
5,631.97.....	5	17,750.86	$\alpha-\phi$	Do.
5,639.74.....	2, h	-----	-----	
5,660.78.....	2, h	17,660.51	-----	
5,691.26.....	2	17,565.96	-----	
5,707.47.....	2	17,516.05	-----	
5,730.46.....	4	17,445.76	-----	Do.
5,774.557.....	3	17,312.54	-----	
5,806.30.....	1, h	17,217.90	-----	
5,813.97.....	0	17,195.19	-----	
5,844.66.....	1, H	17,104.89	-----	
5,910.64.....	1, h, v, +	-----	-----	Do.
5,912.33.....	2	16,909.14	-----	
5,948.37.....	1	16,806.69	-----	Do.
6,005.00.....	4, h, +	-----	-----	
6,079.55.....	3, H, +	-----	-----	Do.
6,081.14.....	0	16,439.74	-----	Do.
6,097.60.....	2	16,395.38	$3d_2-3D_2$	
6,112.70.....	0	16,354.86	-----	
6,129.98.....	3, H, +	-----	-----	
6,178.85.....	1	16,179.76	-----	
6,179.80.....	1	16,177.28	-----	Do.
6,204.51.....	0	16,112.87	-----	
6,214.00.....	2	16,088.24	-----	
6,221.45.....	3	16,068.98	-----	
6,332.220.....	3	15,787.89	-----	
6,345.955.....	2	15,753.71	-----	Do.
6,388.324.....	2	15,649.26	$\xi-\nu_1$	
6,392.175.....	3	15,639.81	-----	
6,405.406.....	2	15,607.49	-----	
6,415.93.....	1, H	15,581.91	$\epsilon-\omega$	
6,454.95.....	2	15,487.72	-----	Do.
6,490.25.....	2	15,403.48	$\alpha-\sigma_1$	
6,503.03.....	1, H	-----	-----	
6,505.404.....	1	15,367.62	-----	
6,517.10.....	2	15,340.01	$\alpha-\sigma_2$	
6,558.150.....	3	15,244.00	-----	Do.
6,564.052.....	3	15,230.30	-----	
6,611.48.....	3	15,121.04	-----	
6,634.63.....	2, H	15,068.28	$\xi-\mu$	
6,640.55.....	1, H	15,054.84	$\beta-\phi$	
6,648.134.....	2	15,037.68	-----	Do.
6,738.99.....	2	14,834.92	-----	
6,778.38.....	2, H, +	-----	-----	
6,806.35.....	2, H, +	-----	-----	
6,811.42.....	1	14,677.18	-----	
6,995.91.....	2, h, +	-----	-----	Do.
7,006.19.....	1, h	14,269.16	-----	
7,032.88.....	1	14,215.00	-----	
7,122.28.....	1	14,036.58	-----	
7,136.29.....	1	14,009.04	$\gamma-\phi$	
7,222.50.....	2, h	13,841.81	-----	Do.
7,362.30.....	1, h	13,578.99	-----	
7,405.50.....	2, h	13,499.76	-----	
7,442.0.....	1, H	13,433.5	-----	
7,592.82.....	1, h	13,166.71	-----	
7,648.28.....	3	13,071.25	-----	Do.
7,755.85.....	2	12,889.95	$\beta-\rho$	
7,764.9.....	2, H	12,874.9	-----	
7,816.9.....	2, H	12,789.2	-----	
7,834.6.....	1, H	12,760.3	-----	

TABLE 7.—Arc Spectrum of Antimony—Continued

λ	Intensity	ν	Classification	Remarks
λ I. A.				
7,837.25.....	2	12,756.08	
7,844.44.....	4	12,744.37	
7,867.17.....	2	12,707.56	$\beta-o_1$	
7,906.73.....	2	12,643.98	$\beta-o_2$	
7,924.65.....	6	12,615.38	
7,933.38.....	1	12,601.50	
7,969.47.....	2, h	12,544.44	
8,062.1.....	2, H	12,400.3	
8,241.74.....	1	12,130.03	
8,257.56.....	2, H	12,106.78	
8,289.0.....	2, H	12,060.8	
8,314.21.....	1	12,024.30	
8,411.65.....	2	11,885.01	$\alpha-u_1$	
8,572.59.....	2	11,661.89	$\gamma-o_1$	
8,619.60.....	2	11,598.29	$\gamma-o_2$	
8,682.88.....	1	11,513.76	
8,700.20.....	1	11,490.83	
8,719.79.....	1	11,465.02	
8,735.0.....	1, H	11,445.0	$\theta-\omega$	
8,789.2.....	1, H	11,374.4	
8,860.3.....	1, H	11,283.2	
8,903.1.....	1, H	11,228.9	
9,019.5.....	1, H	11,084.0	
9,132.30.....	1	10,947.15	
9,519.9.....	20	10,501.5	$\delta-\rho$	From 9519 to 12118 A the intensity numbers are galvanometer deflections.
9,950.5.....	15	10,047.2	
10,079.9.....	35	9,918.0	$\zeta-\phi$	
10,262.9.....	40	9,741.2	$\gamma-v_1$	
10,587.2.....	50	9,442.8	$\epsilon-\rho$	
10,678.0.....	130	9,362.6	
10,742.9.....	50	9,305.9	
10,840.6.....	50	9,222.1	
10,880.3.....	30	9,188.4	$\beta-u_1$	
11,013.4.....	20	9,077.3	
11,082.7.....	15	9,020.6	
11,109.7.....	15	8,998.7	
11,190.3.....	10	8,933.9	
11,268.5.....	45	8,871.9	
11,864.3.....	40	8,426.3	
12,118.9.....	15	8,249.3	

6. BISMUTH

Randall's wave lengths were used from 22554 to 9657 A, where Walters' observations began. Kayser and Runge's numbers extend to 2061 A, while the lines at 2073, 2069, 2049, 2021, and 2020 A are due to Eder and Valenta.²⁵ From 1990 to 1902 A the numbers are those of Takamine and Nitta.²⁶ The region from 1900 to 1400 A has been investigated by L. and E. Bloch²⁷ and by McLennan, Young, and Ireton.²⁸ Since no classifications were obtained in this region we do not give the wave lengths. Tables 8 and 9 contain the term magnitudes, the wave lengths and classi-

²⁵ Atlas Typischer Spectren, 1911.²⁶ Kyoto College of Science Memoirs, 2, p. 117; 1917.²⁷ C. R., 170, p. 320, and 171, p. 709; 1920.²⁸ Proc. Roy. Soc., 98, p. 95; 1920.

fications. The following pairs are characterized by the difference 348.59:

λ	ν
9,657.20	10,352.14
9,342.60	10,700.73
8,761.54	11,410.40
8,501.8	11,758.99

It must be mentioned that we can do away with the orbits $3d_1^A$, $1s$, and ρ , provided we introduce three others at 51031.43 4610.46, and 5103.22. However, this will leave the line $1s-\zeta$ unclassified, and will destroy the symmetry of arrangement of the $3d$ and $3D$ groups. The correct allocation can not be decided without new data.

NOTE ADDED MARCH 20, 1924.—Unfortunately, the wave lengths of Offermann, given in the seventh volume of Kayser and Konen's Handbuch, were not available at the time this paper was written. So far as can be seen, no changes are made necessary by the new data.

TABLE 8.—Term Magnitudes in Bismuth

Name of term	Term magnitude	Differences
σ	4,596.41
ρ	5,015.92
π	5,793.98
ϕ	6,299.53
ξ	6,781.38
ν	8,991.08
μ	11,711.13
λ	11,850.00
κ	13,408.29
ι	14,379.67
θ	15,496.12
η	16,953.70
ζ	18,004.03
ϵ	18,052.13
δ	18,956.53
γ	21,742.76
β	29,703.81
α	30,280.38
$3D_1^A$	41,207.49
$3D_1^B$	41,209.34	1.85
$3D_2$	47,431.76	6,222.42
$1s$	50,944.00
$3d_1^A$	51,435.95	14.05
$3d_1^B$	51,450.00
$3d_2$	62,868.26	11,418.26

TABLE 9.—Arc Spectrum of Bismuth

λ	Intensity	ν	Classification	Remarks
λ vac.				
1,902.6.....	10	52,561	L. and E. Bloch give 1,973.09 and 1,959.57 Å, $\Delta\nu=349.7$.
1,910.2.....	2	52,352	
1,953.9.....	5	51,181	$3d_2-\mu$	
1,959.7.....	2	51,030	$3d_2-\lambda$	
1,973.3.....	2	50,678	
1,988.9.....	4	50,280	
1,990.2.....	1	50,248	
λ I. A.				
2,020.5.....	1	49,476	
2,020.99.....	1	49,464.7	$3d_2-\kappa?$	
2,049.59.....	2	48,774.5	
2,061.71.....	10R	48,487.9	$3d_2-\iota$	
2,068.99.....	2	48,317.4	
2,073.2.....	3	48,218	
2,110.29.....	10R	47,371.9	$3d_2-\theta$	
2,133.66.....	8R	46,853.0	$3d_1^B-\sigma$	
2,134.32.....	10R	46,838.5	$3d_1^A-\sigma$	
2,152.91.....	8R	46,434.0	$3d_1^B-\rho$	
2,153.53.....	4R	46,420.4	$3d_1^A-\rho$	
2,156.96.....	10R	46,346.8	$1s-\sigma$	
2,164.09.....	4R	46,194.2	
2,176.63.....	6R	45,928.2	$1s-\rho$	
2,177.33.....	45,913.4	$3d_2-\eta$	
2,189.63.....	8R	45,655.4	$3d_1^B-\pi$	
2,203.1.....	6U	45,375.70	
2,214.14.....	4	45,150.2	$3d_1^B-\sigma$	
2,224.20.....	2	44,946.0	
2,228.24.....	8R	44,864.6	$3d_2-\xi$	
2,230.63.....	10R	44,816.3	$3d_2-\epsilon$	
2,276.57.....	8R	43,912.1	$3d_2-\delta$	
2,281.32.....	2u	43,820.5	
2,328.19.....	2r	42,938.5	
2,333.79.....	2	42,835.5	$3D_2-\sigma$	
2,354.49.....	2u	42,459.0	$3d_1^B-\nu$	
2,400.90.....	8R	41,638.4	$3D_2-\pi$	
2,430.43.....	2u	41,132.5	$3D_2-\sigma$	
2,433.4.....	4U	41,082.0	Visible at 10.8 volts. May be Sn.
2,448.07.....	4	40,836.2	Visible at 7.8 volts.
2,489.4.....	6U	40,158.0	Not on our plates.
2,515.63.....	6R	39,739.4	$3d_1^B-\mu$	On our plates 2,515 and 2,524 are equally intense.
2,524.49.....	2	39,600.0	$3d_1^B-\lambda$	
2,532.56.....	4U	39,473.9	Not on our plates.
2,582.08.....	2	38,717.0	Do.
2,594.05.....	1	38,538.2	
2,600.64.....	1	38,440.6	$3D_2-\nu$	
2,627.89.....	8R	38,042.0	$3d_1^B-\kappa$	
2,696.59.....	5	37,072.9	
2,696.74.....	6R	37,070.8	$3d_1^B-\iota$	Diffuse in under-water spark.
2,730.35.....	3	36,614.4	$3D_1^B-\sigma$	
2,730.51.....	6R	36,612.3	$3D_1^A-\sigma$	
2,780.47.....	8R	35,954.5	$3d_1^B-\theta$	Raie ultime.
2,798.65.....	4	35,720.1	$3D_2-\mu$	
2,809.63.....	8R	35,581.3	$3D_2-\lambda$	
2,863.75.....	4	34,909.0	$3D_1^A, ^B-\sigma$	
2,883.77.....	1u	34,666.7	
2,897.97.....	10R	34,496.8	$3d_1^B-\eta$	Do.
2,938.30.....	10R	34,023.4	$3D_2-\kappa$	Do.
2,989.03.....	8R	33,445.9	$3d_1^B-\xi$	Do.
2,993.34.....	8R	33,397.7	$3d_1^B-\epsilon$	
3,024.63.....	8R	33,052.3	$3D_2-\iota$	
3,034.87.....	4r	32,940.8	$1s-\xi?$	May be a 14-volt line.
3,067.69.....	10R	32,588.4	$3d_2-\alpha$	Raie ultime.
3,076.61.....	2	32,493.9	$3d_1^B-\delta$	
3,397.17.....	4R	29,427.8	$3D_2-\xi$	Diffuse toward red in under-water spark. Bloch, Jour. de Phys. (6) 3, p. 309, 1922.
3,405.18.....	3	29,359.3	$3D_1^B-\lambda$	
3,405.25.....	2R	29,358.0	$3D_1^A-\lambda$	
3,510.88.....	4R	28,475.0	$3D_2-\delta$	Diffuse toward red in under-water spark.
3,596.11.....	4R	27,799.9	$3D_1^A, ^B-\lambda$	
3,887.90.....	1	25,713.6	$3D_1^B-\theta$	

TABLE 9.—Arc Spectrum of Bismuth—Continued.

λ	Intensity	ν	Classification	Remarks
λ I. A.				
3,888.19.....	1	25, 711. 6	$3D_1^A-\theta$...	Diffuse toward red in under-water spark.
4,121.54.....	6	24, 256. 0	$3D_1^B-\eta$...	
4,121.86.....	6	24, 254. 1	$3D_1^A-\eta$...	Appears at 8.5 volts. Not recorded in arc in air.
4,254.17.....	1u	23, 499. 8	$\alpha-\xi$	
4,272.5.....	2u	23, 398. 7	$3D_1^B-\xi$	Raie ultime.
4,308.18.....	4	23, 205. 2	$3D_1^A-\xi$...	
4,308.54.....	4	23, 203. 2	$3D_1^B-\xi$...	May be a flame line, according to Schwetz.
4,492.62.....	2	22, 252. 5	$3D_1^A-\delta$...	
4,492.99.....	2	22, 250. 7	$3D_1^B-\delta$...	5 A wide.
4,615.09.....	1	21, 662. 0	$3d_2-3D_1^A$	
4,615.53.....	1	21, 659. 9	$3d_2-3D_1^B$	Do.
4,722.54.....	10R	21, 169. 1	$3d_1^B-\alpha$...	
4,733.73.....	4r	21, 119. 1	5 A wide.
5,298.34.....	2r	18, 868. 6	
5,552.35.....	10, B, r	18, 004. 69	Do.
5,599.41.....	3	17, 853. 38	$\beta-\lambda$	
5,742.55.....	6, r	17, 408. 38	5 A wide.
6,134.82.....	6	16, 295. 35	$\beta-\kappa$	
6,184.99.....	2, H	16, 163. 15	Do.
6,364.75.....	1, h	15, 706. 67	
6,475.73.....	3	15, 437. 53	5 A wide.
6,476.24.....	3	15, 436. 32	$3d_2-3D_2$...	
6,991.12.....	4, H	14, 299. 40	Do.
7,036.15.....	2	14, 207. 87	$\beta-\theta$	
7,335.01.....	1	13, 628. 99	5 A wide.
7,441.25.....	1, h	13, 434. 40	
7,502.33.....	2	13, 325. 03	Do.
7,838.70.....	3	12, 753. 25	
7,840.33.....	2	12, 750. 60	$\beta-\eta$	5 A wide.
8,210.83.....	1, h	12, 175. 25	$\delta-\xi$	
8,501.8.....	1, H, b	11, 758. 51	Do.
8,544.54.....	2	11, 699. 70	$\beta-\xi$	
8,579.74.....	1	11, 651. 69	$\beta-\epsilon$	5 A wide.
8,627.9.....	1, H, b	11, 586. 65	
8,754.88.....	2	11, 418. 61	$3d_2-3d_1^B$..	Do.
8,761.54.....	3	11, 409. 93	
8,907.81.....	2	11, 222. 58	$\xi-\xi$	5 A wide.
9,058.62.....	1	11, 035. 74	
9,342.60.....	1, h	10, 700. 28	Do.
9,657.20.....	2	10, 351. 70	
9,828.8.....	1 20 mm	10, 171. 5	$\eta-\xi$	5 A wide.
10,106.1.....	20	9, 892. 3	$\gamma-\lambda$	
10,301.7.....	15	9, 704. 5	Do.
10,540.2.....	8	9, 484. 9	
11,073.2.....	15	9, 028. 4	5 A wide.
11,555.5.....	5	8, 651. 5	
11,711.1.....	160	8, 536. 6	$\alpha-\gamma$	Do.
11,994.5.....	13	8, 334. 9	$\gamma-\kappa$	
12,166.5.....	40	8, 217. 0	5 A wide.
12,296.....	8, 132. 7	
12,690.5.....	30	7, 877. 8	Do.
14,331.5.....	25	6, 975. 8	
22,554.2.....	7	4, 432. 6	5 A wide.
			

¹ From 9828 to 22554 A the intensity numbers represent galvanometer deflections. They are from 5 to 20 times as large as intensities estimated from photographs of the region around 10000 A, when the latter are expressed on a scale of 1 to 10.

7. HIGHER STAGES IN THE BISMUTH SPECTRA

Starting from the fact that the difference 203.65 occurs several times in the 14-volt lines of Bi, it is possible to correlate 13 lines by means of the energy levels given below. These levels are designated by arbitrary letters and the term magnitudes are also arbitrary. It may even occur that the direction of increase of the terms should be reversed. Many spark lines are recorded only

by one observer and frequently the wave lengths are given only to 0.1 Å. For these reasons no attempt was made to push the classifications too far, and a complete list of wave lengths will not be given.

Terms of the 14-Volt Spectrum of Bismuth

Name	ν	$\Delta\nu$
<i>N</i>	78,636.45	1,114.33
<i>M</i>	79,750.78	45.57
<i>L</i>	79,796.35	203.65
<i>K</i>	80,000.00	
<i>J</i>	96,313.20	3,119.23
<i>I</i>	99,432.43	
<i>H</i>	105,993.30	
<i>G</i>	108,933.93	2,940.63

The correlated lines are as follows:

λ I. Å.	Intensity	ν	Classification
3,299.7.....	1u	30,297.4	<i>G-N</i>
3,431.01.....	20	29,137.6	<i>G-L</i>
3,455.2.....	1u	28,934.0	<i>G-K</i>
3,654.4.....	1u	27,356.8	<i>H-N</i>
3,816.1.....	4u	26,197.0	<i>H-L</i>
3,846.0.....	2u	25,993.3	<i>H-K</i>
5,079.52.....	1	19,681.4	<i>I-M</i>
5,091.22.....	1	19,636.2	<i>I-L</i>
5,144.57.....	5	19,432.6	<i>I-K</i>
5,655.58.....	3	17,676.7	<i>J-N</i>
6,036.00.....	2u	16,562.7	<i>J-M</i>
6,052.79.....	1u	16,516.8	<i>J-L</i>
6,128.34.....	3	16,313.1	<i>J-K</i>

A few other frequency differences may be significant. The very intense lines

$$3,792.8 \text{ Å}, \quad 26,357.92 \text{ cm}^{-1}$$

$$4,302.09 \text{ Å}, \quad 23,237.99 \text{ cm}^{-1}$$

appear at 14 volts, and have the frequency difference *I-J*. 3793 Å is quite resistant to self-induction, according to Hemsalech.²⁹ There is a strong pair in the vacuum arc spectrum at 1538.5 and 1533.7 Å, which has the separation *K-L*.


There are two pairs, each including a strong reversed line, which have the separation 300.12:

λ I. Å.	ν
5,818.98 (2u)	17,180.39
5,719.08 (6R)	17,480.50
4,797.67 (3u)	20,837.63
4,729.55 (3R)	21,137.77

The difference 300.12 and the cross difference 3657.26 did not lead to any further classifications.

WASHINGTON, January 20, 1924.

²⁹ Jour. de Phys. (3), 8, p. 562; 1899.



SEP 10 1924

DEPARTMENT OF COMMERCE

BUREAU OF STANDARDS

George K. Burgess, Director

**THEORY OF DETERMINATION OF
ULTRA-RADIO FREQUENCIES BY
STANDING WAVES ON WIRES**

By August Hund

SCIENTIFIC PAPERS OF THE BUREAU OF STANDARDS, No. 491

References back not to be
taken from the literature





